# A high-fat diet impairs reproduction by decreasing the IL1 $\beta$ level in mice treated at immature stage

Jie Zhang<sup>1</sup>, kai Li<sup>2</sup>, Miao Yuan<sup>1</sup>, Jie Zhang<sup>2</sup>, Guizen Huang<sup>1</sup>, Jie Ao<sup>2</sup>, Haoze Tan<sup>1</sup>, Yanyan Li<sup>2</sup>, Di Gong<sup>2</sup>, Jun Li<sup>2</sup>, lei Kang<sup>1</sup>, Nini An<sup>3</sup>, Fei Li<sup>3</sup>, Ping Lin<sup>2</sup>\* and Lugang Huang <sup>1</sup>\*

#### **Supplementary Figure legend**


**Supplemental Figure S1.** Serum testosterone levels of control diet mice (n=10) in different ages.

**Supplemental Figure S2.** Correlation analysis on serum IL1 $\beta$  levels and the mRNA expression of STAR, P450SCC, P45017 $\alpha$ , 3- $\beta$ HSD and 17 $\beta$ -HSD in all experimental mice (n=40).


Supplementary Figure S3. Full-length blots for main figures corresponded to Fig. 3c and 3d.

Supplementary Figure S4. Full-length blots for main figures corresponded to Fig. 5j.

## **Supplementary Figure S1.**



## **Supplementary Figure S2.**



#### **Supplementary Figure S3.**

Figure 3c


NLRP3 ← 130KDa ← 100KDa

Caspase1 ← 55KDa ← 40KDa

Cleaved - Caspase1 ← 15KDa ← 10KDa

Cleaved - Caspase1 ← 10KDa ← 40KDa

Figure 3d



### **Supplementary Figure S4.**

Figure 5j

